



發行人：陳志鴻
 總編輯：周辰熹
 主編：藥品諮詢組
 地址：台南市勝利路138號
 電話：(06)2353535 轉 2515
<http://www.ncku.edu.tw/~pharmacy/>
 八十年十月創刊
 新聞局出版事業登記證
 局版台省誌字第1207號

新藥通知

料號	藥理分類號	藥品名稱
207425	6020	Conjugated estrogens and medroxyprogesterone acetate (Androgens, Estrogens and Progestins) 肝 'Premelle Lite' 0.3mg/1.5mg/tab, 28tab/pack Usual dose: 1 tab/day. 孕婦安全分級：X 備註：本院另備有Premelle 2.5 (0.625/2.5mg/tab, 28tab/pack).
041059	6630	Alanyl Glutamine (Parenteral Nutrition Therapy) 腎 肝 'Dipeptiven' 20gm/100ml/btl Usual dose: IVD, 0.3-0.4 g/kg/day (1.5-2 ml/kg/day). Max. 0.4 g/kg/day (2 ml/kg/day). IVD rate < 0.1 g/kg/hr at max. conc. of 3.5%. 注意事項：1.為高濃度輸注液，不適合直接使用。必須至少以五倍以上可相容溶液之體積混合後靜脈輸注。 2.混合後之滲透壓若大於800 mOsmol/L時，應由中心靜脈輸注。 3.給藥劑量應列入氨基酸總量(最大劑量: 2 g/kg/day)計算，且不應超過氨基酸總量的20%。 4.使用勿超過三週。
400458	4850	Gadobutrol (Magnetic Resonance Radiopaques) 腎 'Gadovist' 604.72mg/ml, 15ml/btl Usual dose: IV. Cranial MRI: 0.1 ml/kg (corresponding to 0.1 mmol/kg). A second dose of 0.1-0.2 ml/kg may be given within 30 min if required. MRA, imaging of 1 field of view: < 75 kg, 7.5 ml; > 75 kg, 10 ml (corresponding to 0.1-0.15 mmol/kg). MRA, imaging > 1 field of view: < 75 kg, 15 ml; > 75 kg, 20 ml (corresponding to 0.2-0.3 mmol/kg).

627128	0020	Nefopam (Non-narcotic Analgesics and Antipyretics) 'Panagesic' 30mg/tab Usual dose: 1 tab qid; single dose may be increased to 90 mg. Max. 300 mg/day. 注意事項：有癲癇病史或正使用MAOI者，不建議使用。
902140	4590	Tretinoin/Fluocinolone/Hydroquinone (Agents for Melasma of Face) 'Tri-Luma' Cream 15g/tube Usual dose: Once daily at night, at least 30 min before bedtime. 注意事項：1.避免曝曬日光。 2.禁忌使用封閉型敷料。 孕婦安全分級：C
101723	5490	Balsalazide (Agents for Ulcerative Colitis) 'Basazyde' 750mg/cap Usual dose: 750 mg tid; Max. 6.75 g/day. 注意事項：禁用於對水楊酸鹽過敏者。 孕婦安全分級：B
262428	0030	Diclofenac (Anti-rheumatics: NSAIDs) 腎 不宜磨 'Voren' EMC 50mg/cap Usual dose: 50 mg bid. 孕婦安全分級：C
681722	5420	Pancrelipase (Digestants and Antiflatulents) 不宜磨 'Protease' E.C 280mg/cap Each cap contains: Lipase.....20000 USP Units Protease..... 75000 USP Units Amylase66400 USP Units Usual dose: Adult & Child > 6 yr: One cap with each meal or snacks. Max. 6000 USP Lipase Units/kg/meal. Child ≤ 6 yr: Individualized by measuring the nitrogen and fat content in the stool. The following are guidelines: Child 6 mo-1 yr: 2000 USP units of lipase with meals; Child 1-6 yr: 4000-8000 USP units of lipase with meals. 注意事項：1.本藥為腸溶微粒膠囊，服用時可將膠囊打開，但不可壓碎或咀嚼。 2.不可與酸鹼值5.5以上之食物混合，以免提早釋放出酵素，造

成口腔黏膜刺激及失去酵素之活性。

孕婦安全分級：C

- | | | |
|--------|------|---|
| 443742 | 6010 | <p>Hydrocortisone (Adrenal Corticosteroids)</p> <p>'Cortema' Enema 1.667mg/ml, 60ml/btl</p> <p>Usual dose: Once daily at bedtime for 21 days. Clinical improvement usually occurs in 3-5 days. For severe cases, treatment may be continued for 2-3 mo. If used for longer than 3 wk, treatment should be gradually tapered over 2-3 wk to alternate day.</p> <p>注意事項：建議使用時側臥，每30分鐘翻身一次，並忍住一小時以上再排出。</p> |
| 828123 | 1240 | <p>Sotalol (Anti-Arrhythmics: Class II) 腎 飯前</p> <p>'Cardol' 160mg/tab</p> <p>Usual dose: 160-240 mg/day in divided doses. Doses up to 640 mg may be necessary in some arrhythmias of ventricular origin.</p> <p>注意事項：支氣管痙攣、高血壓引起之次發性右心室衰竭、明顯的右心室肥大、竇性心跳過慢、二至三度A-V block、休克者禁忌使用。</p> <p>孕婦安全分級：C/B (FDA)</p> |
| 604745 | 5450 | <p>Sodium Phosphate (Laxatives, Cathartics and Bulk-forming Agents) 腎</p> <p>'Fleet Ready to Use Enema' 133ml/btl (Na=37mg/ml)</p> <p>Each ml contains:</p> <p style="padding-left: 40px;">Monobasic sodium phosphate.....160.2 mg</p> <p style="padding-left: 40px;">Dibasic sodium phosphate.....60 mg</p> <p>Usual dose:</p> <p style="padding-left: 40px;">Adult & Child ≥ 12yr: 1 btl for enema use. Max. 1 btl/day.</p> <p>注意事項：1.先天性巨大結腸、腸阻塞、閉鎖性肛門、充血性心臟病衰等疾患勿使用。</p> <p style="padding-left: 40px;">2.腎功能不全、電解質失衡、結腸造口術、使用利尿劑或影響電解質藥品者，應審慎使用。</p> |
| 561751 | 6610 | <p>Magnesium Sulfate Heptahydrate (Replenishers and Regulators of Water and Electrolytes) 腎</p> <p>'Magnesium Sulfate' 100mg/ml, 200ml/btl (Mg= 0.81 mEq/ml)</p> <p>Usual dose: IV or IM, 1-10 g/day.</p> <p style="padding-left: 40px;">Eclampsia: Loading dose, 4 g, then IV 5 ml (0.5 g) every 30 min.</p> <p style="padding-left: 40px;">Magnesium supplement: Mg: 0.81 mEq/ml, for TPN only.</p> <p>注意事項：若用於鎂離子缺乏之補充，僅限用於全靜脈營養輸注液調配使用，勿直接單獨做末梢靜脈點滴注射。</p> |

備註：1.第91次藥委會決議進用，並限婦產科醫師使用。

2.本院另備有Magnesium Sulfate 10%, 20 ml/amp。

孕婦安全分級：A

726940 7290 Polyethylene Glycol/Propylene Glycol (OPH Miscellaneous Agents)

'Systane' Lubricant Eye Drops 15ml/btl

Each ml contains:

Polyethylene glycol 4004.0 mg

Propylene glycol3.0 mg

Usual dose: 1-2 gtt as needed.

注意事項：開封30天後勿繼續使用。

661734 6610 Babyate Oral Solution (Replenishers and Regulators of Water and Electrolytes)

'Babyate' 500ml/btl

Each ml contains:

Sodium Chloride.....2.05 mg

Potassium Citrate Monohydrate.....2.16 mg

Sodium Citrate.....0.98 mg

Dextrose Powder Hydrous.....25.00 mg

Distilled water added to.....1.00 ml

Each liter provides:

Sodium.....45 meq

Potassium.....20 meq

Chloride.....35 meq

Citrate.....30 meq

Dextrose.....25 g

Calories.....100 Kcal/L

Usual dose:

Child< 1 yr: 150 ml/kg/day; 1-4 yr: 125 ml/kg/day; > 4 yr: 100 ml/kg/day.

注意事項：1.出生一週內嬰兒不適用。

2.建議開封後24小時內使用完。

3.僅供口服，不能作為注射使用。

565426 5460 Mebeverine (Anticholinergics and Antispasmodics) 腎 肝 飯前

'Duspatalin' 100mg/SC tab

Usual dose: AC, 100 mg tid-qid.

備註：第91次藥委會決議進用，並刪除 Timepedium 30 mg/cap。