



發行人：陳志鴻
 總編輯：周辰熹
 主編：藥品諮詢組
 地址：台南市勝利路138號
 電話：(06)2353535 轉 2515
<http://www.ncku.edu.tw/~pharmacy/>
 八十年十月創刊
 新聞局出版事業登記證
 局版台省誌字第1207號

新藥通知

料號	藥理分類號	藥品名稱
790132	6310	Rotavirus Vaccine, Live, Oral, Pentavalent (Vaccines and Toxoids) 冷藏 避光 'RotaTeq' 2ml/dose Usual dose: PO. The vaccination course consists of 3 doses. First dose is administered between 6 and 12 wks of age followed by 2 more doses separated at 4-10 wks intervals. Do not administer the 3rd dose after 32 wks of age. 注意事項：1.本品項僅供口服使用，不可注射。 2.不可與任何其他疫苗或溶液混合使用，且勿泡製或稀釋。 備註：本院另備有Rotavirus Vaccine 1 ml/dose.
243625	0610	Desloratadine (Antihistamines) 腎 肝 'Aerius' 5mg/tab Usual dose: AC or with meal. Adult & Child > 12 yr: 5 mg qd. 孕婦安全分級：C
588936	6980	Metoclopramide HCl (Drugs Used in Vertigo and Vomiting) 腎 飯前 'Aswell' Soln 1mg/ml, 60ml/btl Usual dose: 30 min before meals and at bedtime. Gastroesophageal reflux disease: 10-15 ml qid. For intermittent symptoms, single dose may be up to 20 ml. For hypersensitivity patients, 5 ml/dose. Max. recommended therapy is 12 wk. Diabetic gastroparesis: 10 ml qid for 2-8 wk. 孕婦安全分級：B
035554	6340	Adalimumab (Immunomodulators) 冷藏 'Humira' 40mg/pre-filled syringe

Usual dose: SC.

Rheumatoid arthritis, psoriasis with arthropathy and ankylosing spondylitis:

Adult, 40 mg every other week. In rheumatoid arthritis patients without receiving methotrexate, dose may be increase to 40 mg qw.

Crohn's disease: Adult, 160 mg on 1st day (may administer as 4 injections in 1 day or 2 injections daily for 2 consecutive days), followed by 80 mg after 2 wk, then 40 mg q2 wk.

注意事項：不可冷凍保存。

孕婦安全分級：B

820024 6610 Sodium Bicarbonate (Replenishers and Regulators of Water and Electrolytes)
'Sodium Bicarbonate' 600mg/tab (Na: 7.14mEq/tab)

Usual dose: PO, chewed before swallowing.

Adult & Child > 12 yr, 1 tab tid-qid.

Child: 6-12 yr, 1/2 tab tid-qid; 3-6 yr, 1/4 tab tid-qid.

孕婦安全分級：C

606048 1580 Moxifloxacin Ophthalmic Solution (Quinolones)
'Vigamox' 0.5%, 5ml/btl

Usual dose: 1 gtt tid.

注意事項：本品不可為注射之用，不可注射於結膜下及眼睛前房。

孕婦安全分級：C