

成醫藥誌

Pharmacy Forum

發行人：陳 志 鴻
 總編輯：周 辰 熹
 主 編：藥品諮詢組
 地 址：台南市勝利路 138 號
 電 話：(06)2353535 轉 2515
<http://www.ncku.edu.tw/~pharmacy/>
 八十年十月創刊
 新聞局出版事業登記證
 局版台省誌字第1207號

新 藥 通 知

料 號	藥 理	藥品名稱
	分類號	
604351	3320	Mitoxantrone HCl (Anti-neoplastic Agent) 肝 'Mitoxantrone' 20mg/10ml/vial Usual dose: As a single IV dose, 14 mg/M ² repeated at 21-day interval. In combination therapy, dose should be reduced by 2-4 mg/M ² below the dose recommended for single agent usage. 孕婦安全分級：D
126647	7210	Brimonidine tartrate/Timolol Ophthalmic Soln (Drug used to treat Glaucoma) 避光 'Combigan' 2mg/5mg/ml, 5ml/btl Usual dose: 1 gtt bid. 注意事項：使用前應先摘除隱形眼鏡，投藥後15分鐘才可重新配戴。 孕婦安全分級：C 備註：第91次藥委會決議進用，並刪除 Betaxolol Oph Soln 0.5%, 5 ml/btl。
	6610	Peritoneal Dialysis Solution (Replenishers and Regulators of Water and Electrolytes)
681158		Peritoneal Dialysis Solution with 1.5% Glucose, 1.75mmol Calcium 'CAPD/DPCA 2 Peritoneal Dialysis' 2L/bag (Osmolarity: 358mOsmol/L)
681354		Peritoneal Dialysis Solution with 2.3% Glucose, 1.75mmol Calcium 'CAPD/DPCA 4 Peritoneal Dialysis' 2L/bag (Osmolarity: 401mOsmol/L)
681251		Peritoneal Dialysis Solution with 4.25% Glucose, 1.75mmol Calcium 'CAPD/DPCA 3 Peritoneal Dialysis' 2L/bag (Osmolarity: 511mOsmol/L) Each 1 L contains (except glucose): Sodium Chloride.....5.789 g Sodium Lactate.....3.925 g

Calcium Chloride·2H ₂ O.....	0.2573 g
Magnesium Chloride·6H ₂ O.....	0.1017 g
Na ²⁺	134 mmol/L
Ca ²⁺	1.75 mmol/L
Mg ²⁺	0.5 mmol/L
Cl ⁻	103.5 mmol/L
Lactate.....	35 mmol/L
PH.....	5.5

Usual dose: The mode of therapy, frequency of administration, dwell time and exchange volumes required should be specified by the doctor.

- 627128 0020 Nefopam (Non-narcotic Analgesics and Antipyretics)
'Panagesic' 30mg/tab
Usual dose: 1 tab qid; single dose may be increased to 90 mg. Max. 300 mg/day.

注意事項：有癲癇病史或正使用MAOI者，不建議使用。

- 902140 4590 Tretinoin/Fluocinolone/Hydroquinone (Agents for Melasma of Face)
'Tri-Luma' Cream 15g/tube
Usual dose: Once daily at night, at least 30 min before bedtime.
注意事項：1. 避免曝曬日光。
2. 禁忌使用封閉型敷料。

孕婦安全分級：C

- 101723 5490 Balsalazide (Agents for Ulcerative Colitis)
'Basazyde' 750mg/cap
Usual dose: 750 mg tid; Max. 6.75 g/day.
注意事項：禁用於對水楊酸鹽過敏者。

孕婦安全分級：B

- 262428 0030 Diclofenac (Anti-rheumatics: NSAIDs) 腎 不宜磨
'Voren' EMC 50mg/cap
Usual dose: 50 mg bid.

孕婦安全分級：C

- 681722 5420 Pancrelipase (Digestants and Antiflatulents) 不宜磨
'Protease' E.C 280mg/cap
Each cap contains:
Lipase.....20000 USP Units
Protease..... 75000 USP Units

Amylase66400 USP Units

Usual dose:

Adult & Child > 6 yr: One cap with each meal or snacks. Max. 6000 USP

Lipase Units/kg/meal.

Child ≤ 6 yr: Individualized by measuring the nitrogen and fat content in the stool. The following are guidelines: Child 6 mo-1 yr: 2000 USP units of lipase with meals; Child 1-6 yr: 4000-8000 USP units of lipase with meals.

注意事項：1. 本藥為腸溶微粒膠囊，服用時可將膠囊打開，但不可壓碎或咀嚼。

2. 不可與酸鹼值5.5以上之食物混合，以免提早釋放出酵素，造成口腔黏膜刺激及失去酵素之活性。

孕婦安全分級：C

443742 6010 Hydrocortisone (Adrenal Corticosteroids)

'Cortema' Enema 1.667mg/ml, 60ml/btl

Usual dose: Once daily at bedtime for 21 days. Clinical improvement usually occurs in 3-5 days. For severe cases, treatment may be continued for 2-3 mo. If used for longer than 3 wk, treatment should be gradually tapered over 2-3 wk to alternate day.

注意事項：建議使用時側臥，每30分鐘翻身一次，並忍住一小時以上再排出。

828123 1240 Sotalol (Anti-Arrhythmics: Class II) 腎 飯前

'Cardol' 160mg/tab

Usual dose: 160-240 mg/day in divided doses. Doses up to 640 mg may be necessary in some arrhythmias of ventricular origin.

注意事項：支氣管痙攣、高血壓引起之次發性右心室衰竭、明顯的右心室肥大、竇性心跳過慢、二至三度A-V block、休克者禁忌使用。

孕婦安全分級：C/B (FDA)

604745 5450 Sodium Phosphate (Laxatives, Cathartics and Bulk-forming Agents) 腎

'Fleet Ready to Use Enema' 133ml/btl (Na=37mg/ml)

Each ml contains:

Monobasic sodium phosphate.....160.2 mg

Dibasic sodium phosphate.....60 mg

Usual dose:

Adult & Child ≥ 12yr: 1 btl for enema use. Max. 1 btl/day.

注意事項：1. 先天性巨大結腸、腸阻塞、閉鎖性肛門、充血性心臟病衰等疾患勿使用。

2. 腎功能不全、電解質失衡、結腸造口術、使用利尿劑或影響電解質藥品者，應審慎使用。

561751 6610 Magnesium Sulfate Heptahydrate (Replenishers and Regulators of Water and Electrolytes) 腎

'Magnesium Sulfate' 100mg/ml, 200ml/btl (Mg= 0.81 mEq/ml)

Usual dose: IV or IM, 1-10 g/day.

Eclampsia: Loading dose, 4 g, then IV 5 ml (0.5 g) every 30 min.

Magnesium supplement: Mg: 0.81 mEq/ml, for TPN only.

注意事項：若用於鎂離子缺乏之補充，僅限用於全靜脈營養輸注液調配使用，勿直接單獨做末稍靜脈點滴注射。

備註：1. 第91次藥委會決議進用，並限婦產科醫師使用。

2. 本院另備有Magnesium Sulfate 10%, 20 ml/amp。

孕婦安全分級：A

726940 7290 Polyethylene Glycol/Propylene Glycol (OPH Miscellaneous Agents)

'Systane' Lubricant Eye Drops 15ml/btl

Each ml contains:

Polyethylene glycol 4004.0 mg

Propylene glycol3.0 mg

Usual dose: 1-2 gtt as needed.

注意事項：開封30天後勿繼續使用。

661734 6610 Babyate Oral Solution (Replenishers and Regulators of Water and Electrolytes)

'Babyate' 500ml/btl

Each ml contains:

Sodium Chloride.....2.05 mg

Potassium Citrate Monohydrate.....2.16 mg

Sodium Citrate.....0.98 mg

Dextrose Powder Hydrous.....25.00 mg

Distilled water added to.....1.00 ml

Each liter provides:

Sodium.....45 meq

Potassium.....20 meq

Chloride.....35 meq

Citrate.....30 meq

Dextrose.....25 g

Calories.....100 Kcal/L

Usual dose:

Child < 1 yr: 150 ml/kg/day; 1-4 yr: 125 ml/kg/day; > 4 yr: 100 ml/kg/day.

注意事項：1. 出生一週內嬰兒不適用。

2. 建議開封後24小時內使用完。

3. 僅供口服，不能作為注射使用。

565426 5460 Mebeverine (Anticholinergics and Antispasmodics) 腎 肝 飯前

'Duspatalin' 100mg/SC tab

Usual dose: AC, 100 mg tid-qid.

備註：第91次藥委會決議進用，並刪除 Timepedium 30 mg/cap。

465152 6310 Influenza Vaccine (Vaccines and Toxoids) 冷藏

'Fluvirin' 0.5ml/syringe

Usual dose: IM or deep SC.

Adult & Child $\geq$ 3 yr: 0.5 ml.

Child 6-35 mo: 0.25 ml or 0.5 ml.

For child who has not previously been vaccinated, a second dose should be given after an interval of at least 4 wk.

孕婦安全分級：C