



發行人：陳 志 鴻
 總編輯：周 辰 熹
 主 編：藥品諮詢組
 地 址：台南市勝利路 138 號
 電 話：(06)2353535 轉 2515
<http://www.ncku.edu.tw/~pharmacy/>
 八十年十月創刊
 新聞局出版事業登記證
 局版台省誌字第1207號

新 藥 通 知

料 號	藥 理	藥品名稱
	分類號	
681825	7553	Paliperidone ER (Atypical antipsychotics) 腎 不宜磨 'Invega ER' 3mg/tab Usual dose: 3-6 mg once daily, administered in the morning. Dose increases above 6 mg/day should be made only after clinical reassessment and generally should occur at intervals of more than 5 days. When dose increases are indicated, small increments of 3 mg/day are recommended. Max. 12 mg/day. 孕婦安全分級：C
528843	6020	Levonorgestrel (Androgens, Estrogens and Progestins) 'Mirena' 52mg/pack (20mcg/24hr) Usual dose: One unit is inserted into the uterine cavity. One administration is effective for five years. 孕婦安全分級：X
461154	6090	Ibandronate (Hormones Miscellaneous Agents) 腎 'Bonviva' 3mg/3ml/syringe Usual dose: IV only, 3 mg every 3 month. 注意事項：1.僅可靜脈注射，建議以15-30秒的時間靜脈推注。 2.不可與含鈣溶液或其他靜脈注射藥物混合使用。 孕婦安全分級：C
243728	1820	Deferasirox (Metal Antagonists) 腎 肝 飯前 不宜磨 'Exjade' 125mg/tab Usual dose: PO. Adult & Child: Initial, 20 mg/kg once daily; maintenance, adjust in increments of 5-10 mg/kg every 3 to 6 months based on serum ferritin trends; max 30 mg/kg/day.

注意事項：1. 本藥錠加入一杯水、柳橙汁或蘋果汁中攪拌溶解至澄清懸浮液後服用。

2. 本藥錠不可咀嚼或整顆吞服。進食前至少30分鐘空腹服用，因食物會增加藥品之吸收。

347127 6340 Everolimus (Immunomodulators) 肝

'Certican' 0.25mg/tab

Usual dose: Initially, 0.75 mg bid. The dose adjustments based on blood levels achieved (Ctrough: 3-8 ng/ml), tolerability, individual response, change in co-medications and the clinical situation. Dose adjustments can be made at 4-5 days.

注意事項：避免與葡萄柚汁併服。

102651 6340 Basiliximab (Immunomodulators) 冷藏

'Simulect' 20mg/vial

Usual dose: IVD over 20-30 min; regimen should include cyclosporine and corticosteroids.

Prophylaxis of renal transplant rejection:

Adult & Child $\geq$ 35 kg: 20 mg within 2 hr before transplant, then another 20 mg on day 4 after transplantation.

Child < 35 kg: 10 mg within 2 hr before transplant, then another 10 mg on day 4 after transplantation.

孕婦安全分級：B

523729 3390 Lapatinib (Anti-Neoplastic Miscellaneous Agents) 肝 飯前

'Tykerb' 250mg/tab

Usual dose: 1250 mg once daily.

注意事項：LVEF大於及等於grade 3者則應停用。

孕婦安全分級：D