




公告日期	2026/10/14
異動類別	藥品變更
異動原因	包裝變更
異動描述	Acyclovir 400mg/tab (Acylete[®])敵?治廠商來文，自批號D54-0310起，變更標籤製造廠logo、仿單無紙化暨新增電子仿單二維條碼，藥品包裝變更。 ● 舊品用罄，啟用新品。

異動內容		
	舊藥品	新藥品
料號	028420	
學名	Acyclovir	
商品名	Acylete	
中文名	敵?治	
主成份及含量	400mg/tab	
製造廠名稱	中國化學製藥股份有限公司新豐工廠 (中國化學製藥股份有限公司新豐工廠)	
產地	臺灣	
健保代碼	AC39315100	
外觀/圖檔		
用法	口服	

小孩劑量	Note: Obese patients should be dosed using ideal body weight HSV genital infection, first infection: Non-HIV-exposed/-positive: <12 years: 40 - 80 mg/kg/day divided Q8H for 5 - 10 days (Max: 1,000 mg/day) ?12 years: 200 mg 5 times daily or 400 mg TID for 7 - 10 days HIV-exposed/-positive: <12 years: 20 mg/kg/dose TID for 7 to 10 days (Max: 400 mg/dose) ?12 years: 400 mg TID for 5 - 14 days Herpes zoster (shingles), treatment: Children ?12 years and Adolescents: 800 mg 5 times daily for 5 - 7 days Varicella (chickenpox), treatment: ?2 years: 20 mg/kg/dose QID for 5 days. (Max: 3,200 mg/day)
成人劑量	HSV genital infection, initial episode: 400 mg TID or 200 mg 5 times daily for 7-10 days. HSV genital infection, recurrent episode: 400 mg TID for 5 days or 800 mg BID for 5 days or 800 mg TID for 2 days. HSV genital infection, chronic suppressive therapy: 400 mg BID. Herpes zoster (shingles), treatment: 800 mg 5 times daily for 7-10 days Varicella (chickenpox), treatment: 800 mg 5 times daily for 5-7 days
副作用	