




公告日期	2026/2/3
異動類別	藥品變更
異動原因	年度招標
異動描述	Acetylcysteine inj 針 300mg/3mL/amp (Ancare [®])舒得清注射液因年度招標異動，改以Acetylcysteine inj 針 300mg/3mL/amp (Fumucil [®])服優施注射液替代。 舊品用罄，啟用新品。

異動內容		
	舊藥品	新藥品
料號	024250	
學名	Acetylcysteine	
商品名	Ancare	Fumucil
中文名	舒得清注射液	服優施注射液
主成份及含量	300mg/3mL/amp	
製造廠名稱	南光化學製藥股份有限公司 (南光)	瑞士藥廠股份有限公司新市廠 (瑞士藥廠)
產地	臺灣	臺灣
健保代碼	AC57289216	AC22476216
外觀/圖檔	  	  
用法	吸入	

小孩劑量	Infants, Children, and Adolescents: Acetaminophen overdose: Three-bag regimen (21-hour regimen): <u>Loading dose</u>: 150 mg/kg IVD over 1 hour (Max: 15 g) <u>Second dose</u>: 50 mg/kg IVD over 4 hours (Max: 5 g) <u>Third dose</u>: 100 mg/kg IVD over 16 hours (Max: 10 g) Two-bag regimen (20-hour regimen): <u>Loading dose</u>: 200 mg/kg IVD over 4 hour (Max: 20 g) <u>Second dose</u>: 100 mg/kg IVD over 16 hours (Max: 10 g) Mucolytics for respiratory conditions: Inhalation: 1 amp QD-BID
成人劑量	Acetaminophen overdose: Three-bag regimen (21-hour regimen): <u>Loading dose</u>: 150 mg/kg IVD over 1 hour (Max: 15 g) <u>Second dose</u>: 50 mg/kg IVD over 4 hours (Max: 5 g) <u>Third dose</u>: 100 mg/kg IVD over 16 hours (Max: 10 g) Two-bag regimen (20-hour regimen): <u>Loading dose</u>: 200 mg/kg IVD over 4 hour (Max: 20 g) <u>Second dose</u>: 100 mg/kg IVD over 16 hours (Max: 10 g) Mucolytics for respiratory conditions: Inhalation: 1 amp QD-BID
副作用	