



公告日期	2025/7/2
異動類別	藥品變更
異動原因	包裝變更
異動描述	Heparin 25000IU/5mL/vial (Heparin®)肝素鈉注射液自批號AC2105起，瓶蓋由黃色變為藍色。 舊品用罄，啟用新品。

異動內容		
	舊藥品	新藥品
料號	424050	
學名	Heparin	
商品名	Heparin	
中文名	肝素鈉注射液	
主成份及含量	25000IU/5mL/vial	
製造廠名稱	Tai Yu (台裕)	
產地	臺灣	
健保代碼	AC59402240	
外觀/圖檔	 	 
用法	靜脈滴注	
小孩劑量	- Full-dose therapy: Initially 50 U/kg, then IVD 100 U/kg q4h; alternatively 20,000 U/m2/24h. ACCP suggests for neonate or child < 1 yr is 28 U/kg/hr and child > 1 yr is 20 U/kg/hr initially, with dosage adjusted to maintain a target aPTT. - DIC: IV, 25-50 U/kg q4h or by continuous infusion.	

成人劑量	- Full-dose therapy (based on 68 kg adult): Loading dose, 5,000 U IV, then 20,000-40,000 U IVD over 24 hr. Alternatively, in ACCP (American College of Chest Physicians) recommends, LD 80 U/kg then IVD 18 U/kg/hr. Alternatively, LD 10,000 U, then intermittent IV 5,000-10,000 U q4-6h. Alternatively, LD 5,000 U IV, then SC 10,000-20,000 U , followed by 8,000-10,000 U q8h or 15,000-20,000 U q12h. - Acute ischemic complication of AMI: LD 60 U/kg (max. 4000 U), then 12 U/kg/hr (max. 1000 U/hr). Adjust to maintain a target aPTT. - DIC: IV, 50-100 U/kg q4h. If there is no improvement after 4-8 hr, the drug should be discontinued. - DVT prophylaxis: Deep SC, 5,000 U 2 hr prior to surgery and 5,000 U q8-12h post-OP for 7 days or until the patient is fully ambulatory.
副作用	