國立成功大學醫學院附設醫院
National Cheng Kung University Hospital 藥劑部

公告日期	2026/3/25
異動類別	藥品變更
異動原因	包裝變更
異動描述	Rituximab inj 洛希隆 100mg/10mL/vial (Rixathon®) 洛希隆注射劑廠商來文，自批號PP1928起，藥商變更、另增logo、包裝樣式變更為破壞性包裝，外盒增印電子仿單二維條碼，藥品包裝變更。 • 舊品用罄，啟用新品。

異動內容		
	舊藥品	**新藥品**
料號	787852	
學名	Rituximab	
商品名	Rixathon	
中文名	洛希隆注射劑(2支/盒)	
主成份及含量	100mg/10mL/vial	
製造廠名稱	Lek Pharmaceuticals D.D. (Lek)	
產地	斯洛維尼亞	
健保代碼	KC01118229	
外觀/圖檔		
用法	靜脈滴注	
小孩劑量	本品未核准用於兒童青少年。	

成人劑量	**Non-Hodgkin lymphoma:** IVD; 375 mg/m^2 on day 1 of each chemotherapy cycle for up to 8 doses (in combination with chemotherapy) **Rheumatoid arthritis:** IVD; 1000 mg on days 1 and 15 (in combination with MTX), subsequent courses may be administered q24w, but no sooner than q16w **Granulomatosis with polyangiitis (Wegener granulomatosis, GPA):** IVD; Induction: 375 mg/m^2 qw for 4 doses Maintenance: 250 mg q2w for 2 doses, followed by 500 mg q6m, subsequent courses may be administered q24w, but no sooner than q16w **Pemphigus vulgaris:** IVD; 1000 mg q2w for 2 doses (in combination with a tapering course of glucocorticoids), followed by 500 mg at months 12 and every 6 months thereafter or based on clinical evaluation
副作用	