





公告日期	2026/2/9
異動類別	藥品變更
異動原因	年度招標
異動描述	Amikacin inj 250mg/2mL/vial (Aminfec®)安炎注射液因年度招標異動，改以Amikacin inj 500mg/2mL/vial (Acemycin®)時欣黴素注射液替代。 舊品用罄，啟用新品。

異動內容		
	舊藥品	新藥品
料號	061552	061550
學名	Amikacin	
商品名	Aminfec	Acemycin
中文名	安炎注射液	時欣黴素注射液
主成份及含量	250mg/2mL/vial	500mg/2mL/vial
製造廠名稱	生達化學製藥股份有限公司 (生達)	永信藥品工業股份有限公司台中幼獅廠 (永信藥品工業股份有限公司台中幼獅廠)
產地	臺灣	臺灣
健保代碼	AC37550212	AC28477212
外觀/圖檔		
用法	靜脈滴注	

小孩劑量	<i>Please check serum level, and adjust dose accordingly.</i> <i>Avoid peak serum conc > 35 mcg/ml & trough conc > 10 mcg/ml.</i> Neonate: 17.5-18 mg/kg/dose Q24H Infants, Children, and Adolescents: <u>Once-daily dosing</u> (preferred): 15-30 mg/kg/dose IV, IM once daily <u>Conventional dosing</u>: 15-22.5 mg/kg/day divided IV, IM Q8H-Q12H	<i>Please check serum level, and adjust dose accordingly.</i> <i>Avoid peak serum conc > 35 mcg/ml & trough conc > 10 mcg/ml.</i> IM or IVD (> 1-2 hr for Infant) Neonate: 17.5-18 mg/kg/dose Q24H Infants, Children, and Adolescents: <u>Once-daily dosing</u> (preferred): 15-30 mg/kg/dose IV, IM once daily <u>Conventional dosing</u>: 15-22.5 mg/kg/day divided IV, IM Q8H-Q12H
成人劑量	<i>Please check serum level, and adjust dose accordingly.</i> <i>Avoid peak serum conc > 35 mcg/ml & trough conc > 10 mcg/ml.</i> Traditional dosing: 5 mg/kg IM, IV Q8H or 7.5 mg/kg Q12H Once-daily dosing: 15-20 mg/kg IV once daily (Max: 1500 mg/day)	<i>Please check serum level, and adjust dose accordingly.</i> <i>Avoid peak serum conc > 35 mcg/ml & trough conc > 10 mcg/ml.</i> IM or IVD (> 30-60 min) Traditional dosing: 5 mg/kg IM, IV Q8H or 7.5 mg/kg Q12H Once-daily dosing: 15-20 mg/kg IV once daily (Max: 1500 mg/day)
副作用		