




公告日期	2026/2/2
異動類別	藥品變更
異動原因	廠商供藥不及
異動描述	Labetalol inj 針劑 25mg/5mL/amp (Trandate[®]) 湍泰低注射液因廠商無法供應，改以 Labetalol inj 針劑 25mg/5mL/amp (Chenday[®]) 壓泰定注射液替代。 舊品用罄，啟用新品。

異動內容		
	舊藥品	新藥品
料號	522251	
學名	Labetalol	
商品名	Trandate	Chenday
中文名	湍泰低注射液	壓泰定注射液
主成份及含量	25mg/5mL/amp	
製造廠名稱	聯亞藥業股份有限公司新竹廠 (聯亞藥業)	台裕化學製藥廠股份有限公司 (台裕)
產地	臺灣	臺灣
健保代碼	AC32350221	AC49550221
外觀/圖檔		
用法	靜脈滴注	

小孩劑量	Acute severe hypertension: <u>Intermittent IV:</u> Children and Adolescents : 0.2-1 mg/kg/dose (Max: 40 mg/dose) <u>Continuous IV infusion:</u> Infants, Children, and Adolescents : 0.25-3 mg/kg/hour; initiate at lower end of range and titrate up slowly to effect.
成人劑量	Hypertensive emergency: <u>Intermittent IV:</u> Initial: 10-20 mg over 1-2 minutes followed by 20-80 mg every 10 minutes until target blood pressure is reached. Total dose < 200 mg. <u>Continuous IV infusion:</u> Initial loading dose: 10-20 mg over 2 minutes (loading dose is optional if intermittent dose is used prior to continuous infusion), followed by 0.5-2 mg/min (Max: 10 mg/min). Hypertension during anesthesia: Starting dose is 10-20 mg IV following induction of anesthesia. If necessary, additional doses of 5-10 mg were given.
副作用	