



公告日期	2026/10/14
異動類別	藥品變更
異動原因	包裝變更
異動描述	Epoetin β inj 外觀 2000IU/0.3mL/syrg (Recormon®)容可曼針筒裝注射劑廠商來文，自批號30093623起，變更外盒及鋁箔之國際條碼，藥品包裝變更。 舊品用罄，啟用新品。

異動內容		
	舊藥品	新藥品
料號	302855	
學名	Epoetin β	
商品名	Recormon	
中文名	容可曼針筒裝注射劑	
主成份及含量	2000IU/0.3mL/syrg	
製造廠名稱	Roche Diagnostics GmbH (羅氏)	
產地	德國	
健保代碼	KC00876212	
外觀/圖檔		
用法	皮下注射	
小孩劑量	Anemia, CKD related: 1~17 years: <u>Initial:</u> Max: 720 IU/kg IV/SC weekly IV: 40 IU/kg tiw. After 4 wks, may ↑ dose to 80 IU/kg tiw. May further ↑ dose by 20 IU/kg/dose q4w if necessary SC: 20 IU/kg tiw or 60 IU/kg divided into 7 daily doses. May further ↑ dose by 60 IU/kg/week q4w if necessary <u>Maintenance:</u> After reaching Hb target, ↓ weekly dose 50 % of the previously administered dose and titrate q1~2w until stable Hb response is achieved	

成人劑量	Anemia, CKD related: <u>Initial:</u> Max: 720 IU/kg IV/SC weekly /V: 40 IU/kg tiw. After 4 wks, may ↑ dose to 80 IU/kg tiw. May further ↑ dose by 20 IU/kg/dose q4w if necessary SC: 20 IU/kg tiw or 60 IU/kg divided into 7 daily doses. May further ↑ dose by 60 IU/kg/week q4w if necessary <u>Maintenance:</u> After reaching Hb target, ↓ weekly dose 50 % of the previously administered dose and titrate q1~2w until stable Hb response is achieved Anemia, chemotherapy related: SC: <u>Initial:</u> 450 IU/kg qw or divided into tiw or 7 daily doses; Max: 60,000 IU/wk <u>After 4 wks:</u> adjust initial dose based on change in Hb: If Hb ↑ 1~2 g/dL: continue initial dose If Hb ↑ <1 g/dL: consider doubling initial dose (Max: 60,000 IU/wk) If Hb ↑ >2 g/dL: ↓ dose by 25~50 % If Hb >12 g/dL and ?13 g/dL: ↓ dose by 25~50 % If Hb >13 g/dL: hold until Hb ?12 g/dL, reinitiate at ~25% of previous dose <u>Maintenance:</u> After reaching Hb target, ↓ dose by 25~50 % to maintain target Hb level
副作用	