



公告日期	2026/10/16
異動類別	藥品變更
異動原因	到藥通知
異動描述	Penicillin G inj 3MU/vial (Penicillin G Sodium®)苳基青黴素鈉注射劑恢復供應，缺藥期間暫代品項Penicillin G inj 5MU/vial (Penicillin G Sodium®)專案藥品用罄鎖檔停用。 舊品用罄，啟用新品。

異動內容		
	舊藥品	新藥品
料號	685755	685754
學名	Penicillin G	
商品名	Penicillin G Sodium	
中文名	專案藥品	苳基青黴素鈉注射劑
主成份及含量	5MU/vial	3MU/vial
製造廠名稱	General Drugs House Co., LTD. (General Drugs House)	永豐化學工業股份有限公司新莊工廠 (永豐)
產地	泰國	臺灣
健保代碼	X000315221	AC15845216
外觀/圖檔		
用法	靜脈滴注	
小孩劑量	General dose: Neonate: 50,000 units/kg/dose IV, IM Q8H Infants, Children, and Adolescents: 100,000-300,000 units/kg/day IV, IM divided Q4H-Q6H; (Max: 24 MU/day)	Usual dose: [Infants, Children, and Adolescents] Mild to moderate: 100,000-150,000 U/kg/day divided Q6H; Severe infections: 200,000-300,000 U/kg/day divided Q4H-Q6H; [Neonatal] Body weight <1 kg: ???PNA ?14 days: 25,000-50,00

成人劑量	Adults: Bloodstream infection: 12-30 MU/day divided Q4H, <u>adjust the dosage by pathogen.</u> Endocarditis: 12-30 MU/day divided Q4H-Q6H, <u>adjust the dosage by infection focus, pathogen and MIC value.</u> Neurosyphilis: 18-24 MU/day continuous infusion or divided Q4H for 10-14 days.	Usual dose: 3 MU IVD q4-6h Meningitis, Neurosyphilis, Endocarditis: 4 MU IVD q4h
副作用		